

SOME COMPLETELY MONOTONIC FUNCTIONS INVOLVING THE q -TRI-GAMMA AND q -TETRA-GAMMA FUNCTIONS WITH APPLICATIONS

FENG QI

ABSTRACT. Let $\psi_q(x)$ for $q > 0$ stand for the q -psi or q -di-gamma function, the q -analogue of the psi or di-gamma function $\psi(x) = \psi_1(x)$. In the paper, the author proves that the functions $[\psi'_p(x)]^2 + \psi''_p(x)$ for $p \geq 1$ and $[\psi'_q(x) - \ln q]^2 + \psi''_q(x)$ for $0 < q \leq 1$ are completely monotonic with respect to $x \in (0, \infty)$. As applications of these results, the author further establishes monotonic properties of four functions involving the q -di-gamma function $\psi_q(x)$ and two double inequalities for bounding the q -di-gamma function $\psi_q(x)$.

1. INTRODUCTION

1.1. Recall from [20, Chapter XIII] and [31, Chapter IV] that a function f is said to be completely monotonic on an interval I if f has derivatives of all orders on I and

$$0 \leq (-1)^n f^{(n)}(x) < \infty \quad (1.1)$$

for $x \in I$ and $n \geq 0$.

The famous Bernstein-Widder Theorem ([31, p. 161, Theorem 12b]) states that a necessary and sufficient condition that $f(x)$ should be completely monotonic for $0 < x < \infty$ is that

$$f(x) = \int_0^\infty e^{-xt} d\mu(t), \quad (1.2)$$

where $\mu(t)$ is non-decreasing and the integral converges for $0 < x < \infty$. This says that a completely monotonic function $f(x)$ on $(0, \infty)$ is a Laplace transform of the measure $\alpha(t)$.

For more information about properties and applications of completely monotonic functions, please refer to [20, Chapter XIII], [31, Chapter IV], and closely related references therein.

1.2. It is well-known that the classical Euler gamma function $\Gamma(x)$ is defined by

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt, \quad x > 0. \quad (1.3)$$

The logarithmic derivative of $\Gamma(x)$, denoted by $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$, is called psi or di-gamma function, and the derivatives $\psi^{(i)}(x)$ for $i \in \mathbb{N}$ are respectively called

2010 *Mathematics Subject Classification.* Primary 33D05; Secondary 26A48, 26D07, 33B15.

Key words and phrases. completely monotonic function; monotonicity; inequality; q -di-gamma function; q -tri-gamma function; q -tetra-gamma function; application.

This paper was typeset using $\mathcal{A}_{\mathcal{M}}\mathcal{S}\text{-}\mathcal{L}^{\text{A}}\mathcal{T}_{\text{E}}\mathcal{X}$.

polygamma functions. Particularly, the functions $\psi'(x)$ and $\psi''(x)$ are respectively called tri- and tetra-gamma functions, see [1, p. 260] and [10, p. 71].

In [2, p. 208, (4.39)], it was established that

$$[\psi'(x)]^2 + \psi''(x) > \frac{p(x)}{900x^4(x+1)^{10}}, \quad x > 0, \quad (1.4)$$

where

$$\begin{aligned} p(x) = & 75x^{10} + 900x^9 + 4840x^8 + 15370x^7 + 31865x^6 + 45050x^5 \\ & + 44101x^4 + 29700x^3 + 13290x^2 + 3600x + 450. \end{aligned} \quad (1.5)$$

In [32, Theorem 1] and [33, Theorem 2], the functions

$$\frac{x+12}{12x^4(x+1)} - \{[\psi'(x)]^2 + \psi''(x)\}, \quad (1.6)$$

$$[\psi'(x)]^2 + \psi''(x) - \frac{x^2+12}{12x^4(x+1)^2}, \quad (1.7)$$

and

$$[\psi'(x)]^2 + \psi''(x) - \frac{p(x)}{900x^4(x+1)^{10}} \quad (1.8)$$

are respectively proved to be completely monotonic on $(0, \infty)$. From this, we derive

$$\max \left\{ \frac{x^2+12}{12x^4(x+1)^2}, \frac{p(x)}{900x^4(x+1)^{10}} \right\} < [\psi'(x)]^2 + \psi''(x) < \frac{x+12}{12x^4(x+1)} \quad (1.9)$$

for $x > 0$, where $p(x)$ is defined by (1.5). In [13], the function (1.8) was further studied by introducing the notion “completely monotonic degree”.

In [23], it was proved that the function

$$[\psi'(x)]^2 + \psi''(x) - \frac{x^2 + \lambda x + 12}{12x^4(x+1)^2}$$

is completely monotonic on $(0, \infty)$ if and only if $\lambda \leq 0$, so is its negative if and only if $\lambda \geq 4$. Consequently, the double inequality (1.9) was sharpened and refined as

$$\frac{x^2 + \mu x + 12}{12x^4(x+1)^2} < [\psi'(x)]^2 + \psi''(x) < \frac{x^2 + \nu x + 12}{12x^4(x+1)^2} \quad (1.10)$$

on $(0, \infty)$ if and only if $\mu \leq 0$ and $\nu \geq 4$.

It is clear that the inequality

$$\psi''(x) + [\psi'(x)]^2 > 0, \quad x > 0 \quad (1.11)$$

is a weakened version of the inequality (1.4) and the left-hand side inequalities in (1.9) and (1.10). This inequality was deduced or recovered in [7, Theorem 2.1] and [9, Lemma 1.1], employed in [4, Theorem 4.3], [7, Theorem 2.1], [9, Theorem 2.1], [19] and [28, Theorem 2], and generalized in [8, Lemma 1.2 and Remark 1.3]. It was remarked in [29] that a divided difference version of the inequality (1.11) was implicitly obtained in [18].

In [12, 16, 25, 26, 27], among other things, the inequality (1.11) was generalized to the complete monotonicity of the function $\psi''(x) + [\psi'(x)]^2$ and closely related ones.

Recently, some other results in this field were published in [24, 30] and closely related references therein.

1.3. Recall from [5, pp. 493–496] that the q -gamma function, the q -analogue of the gamma function $\Gamma(x)$, is defined for $x > 0$ by

$$\Gamma_q(x) = \begin{cases} (1-q)^{1-x} \prod_{i=0}^{\infty} \frac{1-q^{i+1}}{1-q^{i+x}}, & 0 < q < 1, \\ (q-1)^{1-x} q^{\binom{x}{2}} \prod_{i=0}^{\infty} \frac{1-q^{-(i+1)}}{1-q^{-(i+x)}}, & q > 1, \\ \Gamma(x), & q = 1. \end{cases} \quad (1.12)$$

It has the following basic property

$$\Gamma_q(x) = q^{\binom{x-1}{2}} \Gamma_{1/q}(x). \quad (1.13)$$

The q -psi or q -di-gamma function $\psi_q(x)$, the q -analogue of the psi or di-gamma function $\psi(x)$, is defined for $q > 0$ and $x > 0$ by

$$\psi_q(x) = \begin{cases} \frac{\Gamma'_q(x)}{\Gamma_q(x)}, & q \neq 1, \\ \psi(x), & q = 1. \end{cases} \quad (1.14)$$

Furthermore, the functions $\psi_q^{(k)}(x)$ for $k \in \mathbb{N}$, the q -analogues of the polygamma functions $\psi^{(k)}(x)$, are called q -polygamma functions. In particular, the functions $\psi'_q(x)$ and $\psi''_q(x)$ are called q -tri- and q -tetra-gamma functions.

From (1.12), we obtain for $x \in (0, \infty)$ that

$$\begin{aligned} \psi_q(x) &= -\ln(1-q) + (\ln q) \sum_{k=0}^{\infty} \frac{q^{k+x}}{1-q^{k+x}} \\ &= -\ln(1-q) + (\ln q) \sum_{k=1}^{\infty} \frac{q^{kx}}{1-q^k}, \quad 0 < q < 1 \end{aligned} \quad (1.15)$$

and

$$\begin{aligned} \psi_q(x) &= -\ln(q-1) + (\ln q) \left[x - \frac{1}{2} - \sum_{i=0}^{\infty} \frac{q^{-(i+x)}}{1-q^{-(i+x)}} \right] \\ &= -\ln(q-1) + (\ln q) \left[x - \frac{1}{2} - \sum_{i=1}^{\infty} \frac{q^{-ix}}{1-q^{-i}} \right], \quad q > 1. \end{aligned} \quad (1.16)$$

In [4, pp. 80–81, Lemma 4.6], the inequality (1.11) was generalized as

$$[\psi'_q(x)]^2 + \psi''_q(x) > 0 \quad (1.17)$$

for $q > 1$ and $x > 0$. Applying this inequality, the following conclusion was established in [4, p. 82, Theorem 4.8]: For $q > 1$ and $0 < a < b \leq \infty$, we have

$$\exp\{\alpha[e^{\psi_q(x)}(\psi_q(x)-1)+1]\} \leq \frac{\Gamma_q(x)}{\Gamma_q(x_0)} \leq \exp\{\beta[e^{\psi_q(x)}(\psi_q(x)-1)+1]\} \quad (1.18)$$

on (a, b) , with the best possible constant factors

$$\alpha = \begin{cases} Q_q(b), & \text{if } b < \infty \\ 0, & \text{if } b = \infty \end{cases} \quad (1.19)$$

and $\beta = Q_q(a)$, where $x_0 = x_0(q)$ is the unique positive zero of the function ψ_q for $0 < q \neq 1$, and

$$Q_q(x) = \begin{cases} \frac{\ln \Gamma_q(x) - \ln \Gamma_q(x_0)}{[\psi_q(x) - 1]e^{\psi_q(x)} + 1}, & x \neq x_0, \\ \frac{1}{\psi'_q(x_0)}, & x = x_0. \end{cases} \quad (1.20)$$

For more information on the subject of q -gamma and q -polygamma functions, please refer to [6, 17, 21, 22] and closely related references therein.

1.4. The first aim of this paper is to extend and generalize the inequality (1.17) to the following completely monotonic property.

Theorem 1. *For $q \geq 1$, the function*

$$[\psi'_q(x)]^2 + \psi''_q(x) \quad (1.21)$$

is completely monotonic with respect to $x \in (0, \infty)$.

For $0 < q \leq 1$, the function

$$[\psi'_q(x) - \ln q]^2 + \psi''_q(x) \quad (1.22)$$

is completely monotonic with respect to $x \in (0, \infty)$.

1.5. The second aim of this paper is to apply Theorem 1 to establish the following monotonicity and inequality results.

Theorem 2. *The functions*

$$\phi_q(x) = \begin{cases} \psi_q(x) + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right], & q > 1, \\ \psi(x) + \ln(e^{1/x} - 1), & q = 1, \end{cases} \quad (1.23)$$

$$\varphi_q(x) = \begin{cases} \psi_q(x) + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right), & q > 1, \\ \psi(x) + \ln(e^{1/x} - 1), & q = 1, \end{cases} \quad (1.24)$$

$$\Phi_q(x) = \begin{cases} \psi_q(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right], & 0 < q < 1, \\ \psi(x) + \ln(e^{1/x} - 1), & q = 1, \end{cases} \quad (1.25)$$

$$\Theta_q(x) = \begin{cases} \psi_q(x) - (\ln q)x + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right), & 0 < q < 1, \\ \psi(x) + \ln(e^{1/x} - 1), & q = 1 \end{cases} \quad (1.26)$$

are strictly increasing on $(0, \infty)$. Consequently, we obtain the following inequalities:

(1) *The double inequality*

$$a - \ln(e^{1/x} - 1) < \psi(x) < b - \ln(e^{1/x} - 1) \quad (1.27)$$

holds on $(0, \infty)$ if and only if $a \leq -\gamma$ and $b \geq 0$, where $\gamma = 0.577 \dots$ stands for Euler-Mascheroni's constant.

(2) *For $0 < q < 1$, the double inequality*

$$\begin{aligned} \psi_q(1) + (\ln q)x - \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] &< \psi_q(x) \\ &< \ln \frac{\ln q}{q - 1} + (\ln q)x - \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \end{aligned} \quad (1.28)$$

holds on $(0, \infty)$, where the constants $\psi_q(1)$ and $\ln(\frac{\ln q}{q-1})$ are the best possible.

(3) For $q > 1$, the double inequality

$$\begin{aligned} \psi_q(1) - \ln q - \ln\left(\exp \frac{\ln q}{q^x - 1} - 1\right) &< \psi_q(x) \\ &< \ln \frac{\ln q}{q-1} - \frac{\ln q}{2} - \ln\left(\exp \frac{\ln q}{q^x - 1} - 1\right) \end{aligned} \quad (1.29)$$

holds on $(0, \infty)$, where the scalars $\psi_q(1) - \ln q$ and $\ln \frac{\ln q}{q-1} - \frac{\ln q}{2}$ are the best possible.

2. PROOFS OF THEOREM 1

Now we supply two proofs of Theorem 1.

2.1. First proof of Theorem 1.

2.1.1. For $q = 1$, the function

$$[\psi'_q(x)]^2 + \psi''_q(x) = [\psi'(x)]^2 + \psi''(x), \quad x > 0$$

has been proved to be completely monotonic in [12, 16, 25, 26, 27] and closely-related references therein, as mentioned in the final paragraph of Section 1.2.

2.1.2. For $q > 1$, using the second line in (1.16) gives

$$\frac{[\psi'_q(x)]^2 + \psi''_q(x)}{(\ln q)^2} = 1 + \sum_{i=2}^{\infty} \sum_{j=1}^{i-1} c_j(q, x) c_{i-j}(q, x) - \sum_{i=1}^{\infty} (i-2) c_i(q, x), \quad (2.1)$$

where

$$c_i(q, x) = \frac{i \ln q}{1 - q^{-i}} q^{-ix} > 0.$$

In order to prove the complete monotonicity of the function (1.21), it suffices to show

$$(-1)^k \frac{d^k}{dx^k} \left\{ \frac{[\psi'_q(x)]^2 + \psi''_q(x)}{(\ln q)^2} \right\} \geq 0, \quad k \geq 0. \quad (2.2)$$

For $k = 0$, the inequality (2.2) is equivalent to (1.17). For $k \geq 1$, it is equivalent to

$$(-1)^k \sum_{i=2}^{\infty} \sum_{j=1}^{i-1} \frac{\partial^k [c_j(q, x) c_{i-j}(q, x)]}{\partial x^k} \geq (-1)^k \sum_{i=1}^{\infty} (i-2) \frac{\partial^k [c_i(q, x)]}{\partial x^k}. \quad (2.3)$$

In order to prove (2.3), it suffices to verify

$$(-1)^k \sum_{j=1}^{i-1} \frac{\partial^k [c_j(q, x) c_{i-j}(q, x)]}{\partial x^k} \geq (-1)^k (i-2) \frac{\partial^k [c_i(q, x)]}{\partial x^k}, \quad i \geq 3,$$

that is,

$$\begin{aligned} (-1)^k \sum_{j=1}^{i-1} \frac{(-1)^{k+1} i^k j (i-j) (\ln q)^{k+2} q^{i(1-x)+j}}{(q^j - 1)(q^j - q^i)} \\ \geq (-1)^k (i-2) \frac{(-1)^{k+1} i^{k+1} (\ln q)^{k+1} q^{i(1-x)}}{1 - q^i}, \quad i \geq 3, \end{aligned}$$

which can be further simplified as

$$\sum_{j=1}^{i-1} \frac{j(i-j)}{(q^j-1)(1-q^{i-j})} \leq \frac{(i-2)i}{(\ln q)(1-q^i)}, \quad i \geq 3. \quad (2.4)$$

Since

$$\sum_{j=1}^{i-1} \frac{j(i-j)(q^{i-j}-q^j)}{(q^j-1)(q^{i-j}-1)} = 0,$$

we have

$$\sum_{j=1}^{i-1} \frac{j(i-j)(q^i-1)}{(q^j-1)(q^{i-j}-1)} = \sum_{j=1}^{i-1} \frac{j(i-j)(q^j+1)}{q^j-1}. \quad (2.5)$$

The inequality

$$e^\tau + 1 - \frac{2}{\tau}(e^\tau - 1) = \sum_{k=2}^{\infty} \frac{k-1}{(k+1)!} \tau^k > 0, \quad \tau = j \ln q > 0$$

implies that

$$\frac{j(q^j+1)}{q^j-1} > \frac{2}{\ln q}.$$

This leads to

$$\sum_{j=1}^{i-1} \frac{j(i-j)(q^j+1)}{q^j-1} > \frac{2}{\ln q} \sum_{j=1}^{i-1} (i-j) = \frac{i(i-1)}{\ln q} > \frac{i(i-2)}{\ln q}.$$

Substituting this inequality into (2.5) yields

$$\sum_{j=1}^{i-1} \frac{j(i-j)(q^i-1)}{(q^j-1)(q^{i-j}-1)} > \frac{i(i-2)}{\ln q} \quad (2.6)$$

which is equivalent to the strict inequality in (2.4). Thus, the required result (2.3) is proved. In a word, the function (1.21) is completely monotonic for $q > 1$ on $(0, \infty)$.

2.1.3. For $0 < q < 1$, taking the logarithm and differentiating on both sides of (1.13) reveal

$$\ln \Gamma_q(x) = (\ln q) \binom{x-1}{2} + \ln \Gamma_{1/q}(x), \quad (2.7)$$

$$\psi_q(x) = (\ln q) \left(x - \frac{3}{2}\right) + \psi_{1/q}(x), \quad (2.8)$$

$$\psi'_q(x) = \ln q + \psi'_{1/q}(x), \quad (2.9)$$

$$\psi_q^{(k)}(x) = \psi_{1/q}^{(k)}(x), \quad k \geq 2. \quad (2.10)$$

Therefore, when $0 < q \leq 1$, we have

$$[\psi'_{1/q}(x)]^2 + \psi''_{1/q}(x) = [\psi'_q(x) - \ln q]^2 + \psi''_q(x)$$

on $(0, \infty)$. As a result, the function (1.22) is completely monotonic for $0 < q < 1$ on $(0, \infty)$. The first proof of Theorem 1 is complete.

2.2. Second proof of Theorem 1. By the noted Bernstein-Widder Theorem ([31, p. 161, Theorem 12b]) mentioned in Section 1.1, it is sufficient to write the function (1.21) as a Laplace transform of a measure μ and then verify that the measure is positive. In fact, after some elementary computations necessary for rewriting the expression

$$\frac{[\psi'_q(x)]^2 + \psi''_q(x)}{(\ln q)^2} \quad (2.11)$$

by virtue of the Stieltjes integral representation

$$\psi_q(x) = -\ln(1-q) - \int_0^\infty \frac{e^{-xt}}{1-e^{-t}} d\gamma_q(t) \quad (2.12)$$

for $0 < q < 1$ and $x > 0$, which was established in [17], where

$$\gamma_q(t) = -\ln q \sum_{k=1}^{\infty} \delta(t + k \ln q), \quad (2.13)$$

the measure is found to be

$$\begin{aligned} \mu &= \delta(0) + \frac{\ln q}{1-q^{-1}} \delta(\ln q) \\ &+ (\ln q)^2 \sum_{i=2}^{\infty} \left[\sum_{j=1}^{i-1} \frac{j(i-j)}{(1-q^{-i})(1-q^{j-i})} - (i-2) \frac{i \ln q}{1-q^{-i}} \right] \delta(i \ln q). \end{aligned} \quad (2.14)$$

The computation in Section 2.1.2 shows that this is a positive measure. The second proof of Theorem 1 is sketched.

3. PROOF OF THEOREM 2

3.1. For $q = 1$, the increasingly monotonicity of the function

$$\psi(x) + \ln(e^{1/x} - 1) \quad (3.1)$$

on $(0, \infty)$ was proved in [3, pp. 387–388] and [15, 19] respectively by different approaches.

3.2. In [14, p. 1245, Theorem 4.4], it was deduced that

$$\psi_q^{(k-1)}(x+1) - \psi_q^{(k-1)}(x) = -(\ln q) \frac{d^{k-1}}{dx^{k-1}} \left(\frac{q^x}{1-q^x} \right), \quad 0 < q < 1, \quad k \in \mathbb{N}. \quad (3.2)$$

Substituting (2.9) and (2.10) into (3.2), we easily procure that

$$\psi_{1/q}^{(k-1)}(x+1) - \psi_{1/q}^{(k-1)}(x) = -(\ln q) \frac{d^{k-1}}{dx^{k-1}} \left(\frac{q^x}{1-q^x} \right)$$

for $0 < q < 1$, $k \geq 2$, and $x > 0$. Replacing q by $\frac{1}{q}$ in above identity leads to

$$\psi_q^{(k-1)}(x+1) - \psi_q^{(k-1)}(x) = (\ln q) \frac{d^{k-1}}{dx^{k-1}} \left(\frac{1}{q^x - 1} \right), \quad q > 1, \quad k \geq 2. \quad (3.3)$$

Similarly, combining (2.8) with (3.2), we obtain

$$\psi_q(x+1) - \psi_q(x) = \frac{(\ln q)q^x}{q^x - 1}, \quad q > 1, \quad x > 0. \quad (3.4)$$

For $0 < q \neq 1$, we have

$$\begin{aligned}\exp[\phi_q(x)] &= \exp[\psi_q(x)] \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \\ &= \exp \left[\psi_q(x) + \frac{(\ln q)q^x}{q^x - 1} \right] - \exp[\psi_q(x)] \\ &= \exp[\psi_q(x+1)] - \exp[\psi_q(x)],\end{aligned}$$

where the equality in the final line is obtained by using (3.2) and (3.4). Further differentiating gives

$$\begin{aligned}\frac{d \exp[\phi_q(x)]}{dx} &= \psi'_q(x+1) \exp[\psi_q(x+1)] - \psi'_q(x) \exp[\psi_q(x)], \\ \frac{d\{\psi'_q(x) \exp[\psi_q(x)]\}}{dx} &= \{\psi''_q(x) + [\psi'_q(x)]^2\} \exp[\psi_q(x)].\end{aligned}$$

The complete monotonicity of the function (1.21) in Theorem 1 implies the positivity of the function (1.21) for $q > 1$ on $(0, \infty)$, that is, the inequality (1.17). Hence, the function $\psi'_q(x) \exp[\psi_q(x)]$ is increasing, and so $\frac{d \exp[\phi_q(x)]}{dx} > 0$ for $q > 1$ on $(0, \infty)$. As a result, the function $\exp[\phi_q(x)]$, and thus $\phi_q(x)$, is increasing for $q > 1$ on $(0, \infty)$.

For $0 < q \neq 1$, we also have

$$\begin{aligned}\exp[\Phi_q(x)] &= \exp[\psi_q(x) - (\ln q)x] \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \\ &= \exp \left[\psi_q(x) + \frac{(\ln q)q^x}{q^x - 1} - (\ln q)x \right] - \exp[\psi_q(x) - (\ln q)x] \\ &= \exp[\psi_q(x+1) - (\ln q)x] - \exp[\psi_q(x) - (\ln q)x],\end{aligned}$$

where the equality in the final line is deduced by using (3.2) and (3.4). Further computation gives

$$\begin{aligned}\frac{d \exp[\Phi_q(x)]}{dx} &= [\psi'_q(x+1) - \ln q] \exp[\psi_q(x+1) - (\ln q)x] \\ &\quad - [\psi'_q(x) - \ln q] \exp[\psi_q(x) - (\ln q)x]\end{aligned}$$

and

$$\begin{aligned}\frac{d\{[\psi'_q(x) - \ln q] \exp[\psi_q(x) - (\ln q)x]\}}{dx} \\ = \{\psi''_q(x) + [\psi'_q(x) - \ln q]^2\} \exp[\psi_q(x) - (\ln q)x].\end{aligned}$$

The complete monotonicity of the function (1.22) in Theorem 1 implies the positivity of the function (1.22) for $0 < q < 1$ on $(0, \infty)$. Hence, the function

$$\{[\psi'_q(x) - \ln q] \exp[\psi_q(x) - (\ln q)x]\}$$

is increasing, and so $\frac{d \exp[\Phi_q(x)]}{dx} > 0$ for $0 < q < 1$ on $(0, \infty)$. As a result, the function $\exp[\Phi_q(x)]$, and thus $\Phi_q(x)$, is increasing for $0 < q < 1$ on $(0, \infty)$.

For $0 < q < 1$, utilizing (2.8) yields

$$\begin{aligned}\Phi_q(x) &= (\ln q) \left(x - \frac{3}{2} \right) + \psi_{1/q}(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \\ &= \psi_{1/q}(x) + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] - \frac{3}{2} \ln q \\ &= \psi_{1/q}(x) + \ln \left[\exp \frac{\ln(1/q)}{(1/q)^x - 1} - 1 \right] + \frac{3}{2} \ln(1/q).\end{aligned}\tag{3.5}$$

Replacing $\frac{1}{q}$ by q in above equation leads to the increasingly monotonicity of the function $\varphi_q(x)$ for $q > 1$ on $(0, \infty)$.

For $q > 1$, employing (2.8) gives

$$\begin{aligned}\phi_q(x) &= \psi_{1/q}(x) - \left(\ln \frac{1}{q} \right) \left(x - \frac{3}{2} \right) + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \\ &= \psi_{1/q}(x) - \left(\ln \frac{1}{q} \right) x + \ln \left[\exp \frac{\ln(1/q)}{(1/q)^x - 1} - 1 \right] + \frac{3}{2} \ln \frac{1}{q}.\end{aligned}\tag{3.6}$$

Substituting $\frac{1}{q}$ by q in above equation reduces to the increasingly monotonicity of the function $\Theta_q(x)$ for $0 < q < 1$ on $(0, \infty)$.

3.3. The double inequality (1.27) was sharpened in [8, Theorem 2.8]. It can also be derived from the increasing monotonicity of the function (3.1) on $(0, \infty)$ and the limits

$$\lim_{x \rightarrow 0^+} [\psi(x) + \ln(e^{1/x} - 1)] = -\gamma\tag{3.7}$$

and

$$\lim_{x \rightarrow \infty} [\psi(x) + \ln(e^{1/x} - 1)] = 0.\tag{3.8}$$

For details about calculating these two limits, please refer to [15, Theorem 1].

From (1.15), it is easy to see that

$$\lim_{x \rightarrow \infty} \psi_q(x) = -\ln(1 - q), \quad q \in (0, 1).\tag{3.9}$$

In addition, for $q \in (0, 1)$, we have

$$\lim_{x \rightarrow \infty} \left\{ \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] - (\ln q)x \right\} = \lim_{x \rightarrow \infty} \ln \left[\frac{\exp \frac{(\ln q)q^x}{q^x - 1} - 1}{q^x} \right] = \ln \left(\ln \frac{1}{q} \right).$$

Combining them with the increasing monotonicity of $\Phi_q(x)$, we acquire

$$\psi_q(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] < \ln \left(\ln \frac{1}{q} \right) - \ln(1 - q) = \ln \left(\frac{\ln q}{q - 1} \right)$$

for $q \in (0, 1)$ on $(0, \infty)$, which is equivalent to the right-hand side inequality in (1.28).

From (3.2) for $k = 1$ and the limit

$$\lim_{x \rightarrow 0^+} \psi_q(x) = -\infty\tag{3.10}$$

for $0 < q < 1$, we obtain

$$\begin{aligned}
\psi_q(x) + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] &= \ln \exp[\psi_q(x)] + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \\
&= \ln \left\{ \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] \exp[\psi_q(x)] \right\} \\
&= \ln \left\{ \exp \left[\frac{(\ln q)q^x}{q^x - 1} + \psi_q(x) \right] - \exp[\psi_q(x)] \right\} \\
&= \ln \{ \exp[\psi_q(x+1)] - \exp[\psi_q(x)] \} \\
&\rightarrow \psi_q(1)
\end{aligned} \tag{3.11}$$

as $x \rightarrow 0^+$, so $\lim_{x \rightarrow 0^+} \Phi_q(x) = \psi_q(1)$ for $0 < q < 1$. Combining this with the increasing monotonicity of $\Phi_q(x)$, we acquire $\Phi_q(x) > \psi_q(1)$ for $0 < q < 1$ on $(0, \infty)$, that is,

$$\psi_q(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] > \psi_q(1), \quad 0 < q < 1, \quad x \in (0, \infty).$$

The required inequality in the left-hand side of (1.28) is proved.

For $0 < q < 1$, by using (3.2) for $k = 1$, we have

$$\begin{aligned}
\Theta_q(x) &= \psi_q(x) + \frac{(\ln q)q^x}{q^x - 1} - \frac{(\ln q)q^x}{q^x - 1} - (\ln q)x + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right) \\
&= \psi_q(x+1) + \ln \frac{\exp[\ln q/(q^x - 1)] - 1}{q^x \exp[(\ln q)q^x/(q^x - 1)]} \\
&\rightarrow \psi_q(1) - \ln q
\end{aligned} \tag{3.12}$$

as $x \rightarrow 0^+$. Since $\Theta_q(x)$ is strictly increasing on $(0, \infty)$, it follows that

$$\psi_q(x) - (\ln q)x + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right) > \psi_q(1) - \ln q$$

for $0 < q < 1$ on $(0, \infty)$, which is included in the left-hand side inequality in (1.28).

For $q > 1$, by a similar argument as in (3.12), we have

$$\varphi_q(x) = \psi_q(x+1) + \ln \frac{\exp[\ln q/(q^x - 1)] - 1}{\exp[(\ln q)q^x/(q^x - 1)]} \rightarrow \psi_q(1) - \ln q$$

as $x \rightarrow 0^+$. Furthermore, from (2.8) and (3.9), it follows that

$$\begin{aligned}
\varphi_q(x) &= \psi_{1/q}(x) + (\ln q) \left(x - \frac{3}{2} \right) + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right) \\
&= \psi_{1/q}(x) - \frac{3 \ln q}{2} + \ln \left[q^x \left(\exp \frac{\ln q}{q^x - 1} - 1 \right) \right] \\
&\rightarrow -\ln \left(1 - \frac{1}{q} \right) - \frac{3 \ln q}{2} + \ln(\ln q) \\
&= \ln \frac{\ln q}{q - 1} - \frac{\ln q}{2}
\end{aligned}$$

as $x \rightarrow \infty$. Hence, we have

$$\psi_q(1) - \ln q < \varphi_q(x) < \ln \frac{\ln q}{q - 1} - \frac{\ln q}{2}$$

for $q > 1$ on $(0, \infty)$. The required double inequality (1.29) is proved.

For $q > 1$, the limits (3.10) and (3.11) are still valid, then $\lim_{x \rightarrow 0^+} \phi_q(x) = \psi_q(1)$ for $q > 1$, and so $\phi_q(x) > \psi_q(1)$ which is contained in the left-hand side inequality in (1.29). The proof of Theorem 2 is complete.

4. REMARKS

Finally we remark something else.

Remark 1. From Theorem 1, as in the proof of Theorem 2, it is obtained that the functions

$$\psi'_q(x)e^{\psi_q(x)}, \quad q \geq 1 \quad (4.1)$$

and

$$[\psi'_q(x) - \ln q]e^{\psi_q(x) - (\ln q)x}, \quad 0 < q \leq 1 \quad (4.2)$$

are strictly increasing on $(0, \infty)$.

In [9, Lemma 1.2] and [11, pp. 241–242], it was obtained that the function $\psi'(x)e^{\psi(x)}$ is strictly increasing and less than 1 on $(0, \infty)$. So we naturally ask a question: Can one bound the functions (4.1) and (4.2) for $0 < q \neq 1$?

Remark 2. The formulas (3.2), (3.3), and (3.4) together are q -analogues of the recursion formula

$$\psi^{(n-1)}(x+1) = \psi^{(n-1)}(x) + (-1)^{n-1} \frac{(n-1)!}{x^n} \quad (4.3)$$

for $n \in \mathbb{N}$ on $(0, \infty)$, see [1, pp. 258 and 260, 6.3.5 and 6.4.6].

Moreover, it is easy to see that for $x > 0$ the q -analogue of the recursion formula

$$\Gamma(x+1) = x\Gamma(x) \quad (4.4)$$

is

$$\Gamma_q(x+1) = \frac{1-q^x}{1-q} \Gamma_q(x), \quad q \neq 1. \quad (4.5)$$

Remark 3. We remark that the recursion formulas (3.2), (3.3), and (3.4) may be used to provide an alternative proof of the complete monotonicity of the functions (1.21) and (1.22) along with the similar way as in [12, 25, 26, 27, 32, 33] and closely related references therein. We will carry out this work in a subsequent paper.

Remark 4. As $q \rightarrow 1^-$ or $q \rightarrow 1^+$, the double inequalities (1.28) and (1.29) become (1.27), so Theorem 2 generalizes some results obtained in [3, 8, 15, 19] and closely related references therein to q -analogues.

Remark 5. The strict concavity of the function (3.1) was presented and generalized in [15]. So we conjecture that the four functions $\phi_q(x)$, $\varphi_q(x)$, $\Phi_q(x)$, and $\Theta_q(x)$ in Theorem 2 may also be strictly concave on $(0, \infty)$.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 9th printing, Washington, 1970.
- [2] H. Alzer, *Sharp inequalities for the digamma and polygamma functions*, Forum Math. **16** (2004), no. 2, 181–221; Available online at <http://dx.doi.org/10.1515/form.2004.009>.
- [3] H. Alzer, *Sharp inequalities for the harmonic numbers*, Expo. Math. **24** (2006), no. 4, 385–388.

- [4] H. Alzer and A. Z. Grinshpan, *Inequalities for the gamma and q -gamma functions*, J. Approx. Theory **144** (2007), 67–83; Available online at <http://dx.doi.org/10.1016/j.jat.2006.04.008>.
- [5] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999.
- [6] R. Askey, *The q -gamma and q -beta functions*, Appl. Anal. **8** (1978), 125–141.
- [7] N. Batir, *An interesting double inequality for Euler's gamma function*, J. Inequal. Pure Appl. Math. **5** (2004), no. 4, Art. 97; Available online at <http://www.emis.de/journals/JIPAM/article452.html>.
- [8] N. Batir, *On some properties of digamma and polygamma functions*, J. Math. Anal. Appl. **328** (2007), no. 1, 452–465; Available online at <http://dx.doi.org/10.1016/j.jmaa.2006.05.065>.
- [9] N. Batir, *Some new inequalities for gamma and polygamma functions*, J. Inequal. Pure Appl. Math. **6** (2005), no. 4, Art. 103; Available online at <http://www.emis.de/journals/JIPAM/article577.html>.
- [10] P. S. Bullen, *A Dictionary of Inequalities*, Addison Wesley Longman Limited, Harlow, 1998.
- [11] N. Elezović, C. Giordano, and J. Pečarić, *The best bounds in Gautschi's inequality*, Math. Inequal. Appl. **3** (2000), 239–252.
- [12] B.-N. Guo and F. Qi, *A class of completely monotonic functions involving divided differences of the psi and tri-gamma functions and some applications*, J. Korean Math. Soc. **48** (2011), no. 3, 655–667; Available online at <http://dx.doi.org/10.4134/JKMS.2011.48.3.655>.
- [13] B.-N. Guo and F. Qi, *A completely monotonic function involving the tri-gamma function and with degree one*, Appl. Math. Comput. **218** (2012), no. 19, 9890–9897; Available online at <http://dx.doi.org/10.1016/j.amc.2012.03.075>.
- [14] B.-N. Guo and F. Qi, *Properties and applications of a function involving exponential functions*, Commun. Pure Appl. Anal. **8** (2009), no. 4, 1231–1249; Available online at <http://dx.doi.org/10.3934/cpaa.2009.8.1231>.
- [15] B.-N. Guo and F. Qi, *Some properties of the psi and polygamma functions*, Hacet. J. Math. Stat. **39** (2010), no. 2, 219–231.
- [16] B.-N. Guo, F. Qi, and H. M. Srivastava, *Some uniqueness results for the non-trivially complete monotonicity of a class of functions involving the polygamma and related functions*, Integral Transforms Spec. Funct. **21** (2010), no. 11, 103–111; Available online at <http://dx.doi.org/10.1080/10652461003748112>.
- [17] M. E. H. Ismail and M. E. Muldoon, *Inequalities and monotonicity properties for gamma and q -gamma functions*, in: R.V.M. Zahar (Ed.), Approximation and Computation: A Festschrift in Honour of Walter Gautschi, ISNM, Vol. **119**, Birkhäuser, Basel, 1994, 309–323.
- [18] D. K. Kazarinoff, *On Wallis' formula*, Edinburgh Math. Notes **1956** (1956), no. 40, 19–21.
- [19] W.-H. Li, F. Qi, and B.-N. Guo, *On proofs for monotonicity of a function involving the psi and exponential functions*, Analysis (Munich) **33** (2013), no. 1, in press; Available online at <http://dx.doi.org/10.1524/anly.2012.1175>.
- [20] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer, Dordrecht, 1993.
- [21] D. S. Moak, *The q -gamma function for $q > 1$* , Aequationes Math. **20** (1980), 278–285.
- [22] D. S. Moak, *The q -gamma function for $x < 0$* , Aequationes Math. **21** (1980), 179–191.
- [23] F. Qi, *Complete monotonicity of a family of functions involving the tri- and tetra-gamma functions*, submitted.
- [24] F. Qi, P. Cerone, and S. S. Dragomir, *Complete monotonicity of a function involving the divided difference of psi functions*, Bull. Aust. Math. Soc. (2013), in press; Available online at <http://dx.doi.org/10.1017/S0004972712001025>.
- [25] F. Qi and B.-N. Guo, *Completely monotonic functions involving divided differences of the di- and tri-gamma functions and some applications*, Commun. Pure Appl. Anal. **8** (2009), no. 6, 1975–1989; Available online at <http://dx.doi.org/10.3934/cpaa.2009.8.1975>.
- [26] F. Qi and B.-N. Guo, *Necessary and sufficient conditions for a function involving divided differences of the di- and tri-gamma functions to be completely monotonic*, Available online at <http://arxiv.org/abs/0903.3071>.
- [27] F. Qi and B.-N. Guo, *Necessary and sufficient conditions for functions involving the tri- and tetra-gamma functions to be completely monotonic*, Adv. Appl. Math. **44** (2010), no. 1, 71–83; Available online at <http://dx.doi.org/10.1016/j.aam.2009.03.003>.

- [28] F. Qi and B.-N. Guo, *Sharp inequalities for the psi function and harmonic numbers*, Available online at <http://arxiv.org/abs/0902.2524>.
- [29] F. Qi and Q.-M. Luo, *Bounds for the ratio of two gamma functions—From Wendel's and related inequalities to logarithmically completely monotonic functions*, Banach J. Math. Anal. **6** (2012), no. 2, 132–158.
- [30] F. Qi, Q.-M. Luo, and B.-N. Guo, *Complete monotonicity of a function involving the divided difference of digamma functions*, Sci. China Math. (2013), in press.
- [31] D. V. Widder, *The Laplace Transform*, Princeton University Press, 1946.
- [32] J.-L. Zhao, B.-N. Guo, and F. Qi, *A completely monotonic function involving the tri- and tetra-gamma functions*, Math. Slovaca **63** (2013), no. 1-2, in press.
- [33] J.-L. Zhao, B.-N. Guo, and F. Qi, *Complete monotonicity of two functions involving the tri- and tetra-gamma functions*, Period. Math. Hungar. **65** (2012), no. 1, 147–155; Available online at <http://dx.doi.org/10.1007/s10998-012-9562-x>.

SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

URL: <http://qifeng618.wordpress.com>